
Contents

| | |
|---|-----|
| PREFACE | xii |
| NOTATIONS AND SYMBOLS | xv |
| CHAPTER 1. PURE LIQUIDS | 1 |
| 1.1 Macroscopic modeling of liquids | 1 |
| 1.2. Distribution of molecules in a liquid | 2 |
| 1.2.1. Molecular structure of a non- associated liquid | 3 |
| 1.2.2. The radial distribution function | 4 |
| 1.2.3 The curve representative of the radial distribution function | 6 |
| 1.2.4. Calculation of the macroscopic thermodynamic values | 8 |
| 1.3. Models extrapolated from gases or solids | 9 |
| 1.3.1. Guggenheim's smoothed potential model | 10 |
| 1.3.2. Mie's harmonic oscillator model | 13 |
| 1.3.3. Determination of the free volume on the basis of the dilation and the compressibility | 15 |
| 1.4. Lennard-Jones and Devonshire cellular model | 16 |
| 1.5. Cellular and vacancies model | 25 |
| 1.6. Eyring's semi-microscopic formulation of the vacancy model | 29 |
| 1.7. Comparison between the different microscopic models and experimental results | 32 |

| | |
|---|-----------|
| CHAPTER 2. MACROSCOPIC MODELING OF LIQUID MOLECULAR SOLUTIONS | 37 |
| 2.1. Macroscopic modeling of the Margules expansion | 38 |
| 2.2. General representation of a solution with several components. | 39 |
| 2.3. Macroscopic modeling of the Wagner expansions | 40 |
| 2.3.1. Definition of the Wagner interaction coefficients | 40 |
| 2.3.2. Example of a ternary solution: experimental determination of Wagner's interaction coefficients. | 41 |
| 2.4. Dilute ideal solutions | 43 |
| 2.4.1. Thermodynamic definition of a dilute ideal solution | 43 |
| 2.4.2. Activity coefficients of a component with a pure-substance reference | 44 |
| 2.4.3. Excess Gibbs energy of an ideal dilute solution | 44 |
| 2.4.4. Enthalpy of mixing for an ideal dilute solution | 45 |
| 2.4.5. Excess entropy of a dilute ideal solution | 46 |
| 2.4.6. Molar heat capacity of an ideal dilute solution at constant pressure | 46 |
| 2.5. Associated solutions. | 46 |
| 2.5.1. Example of the study of an associated solution | 47 |
| 2.5.2. Relations between the chemical potentials of the associated solution. | 49 |
| 2.5.3. Calculating the extent of the equilibrium in an associated solution | 50 |
| 2.5.4. Calculating the activity coefficients in an associated solution | 50 |
| 2.5.5. Definition of a regular solution. | 51 |
| 2.5.6. Strictly-regular solutions | 52 |
| 2.5.7. Macroscopic modeling of strictly-regular binary solutions. | 53 |
| 2.5.8. Extension of the model of a strictly-regular solution to solutions with more than two components | 56 |
| 2.6. Athermic solutions | 57 |
| 2.6.1. Thermodynamic definition of an athermic solution | 58 |
| 2.6.2. Variation of the activity coefficients with temperature in an athermic solution. | 58 |

| | |
|---|------------|
| 2.6.3. Molar entropy and Gibbs energy of mixing for an athermic solution | 58 |
| 2.6.4. Molar heat capacity of an athermic solution | 59 |
| CHAPTER 3. MICROSCOPIC MODELING OF LIQUID MOLECULAR SOLUTIONS | 61 |
| 3.1. Models of binary solutions with molecules of similar dimensions | 62 |
| 3.1.1. The microscopic model of a perfect solution | 68 |
| 3.1.2. Microscopic description of strictly-regular solutions | 70 |
| 3.1.3. Microscopic modeling of an ideal dilute solution | 72 |
| 3.2. The concept of local composition | 74 |
| 3.2.1. The concept of local composition in a solution | 74 |
| 3.2.2. Energy balance of the mixture | 76 |
| 3.2.3. Warren and Cowley's order parameter | 78 |
| 3.2.4. Model of Fowler & Guggenheim's quasi-chemical solution | 80 |
| 3.3. The quasi-chemical method of modeling solutions | 87 |
| 3.4. Difference of the molar volumes: the combination term | 92 |
| 3.4.1. Combinatorial excess entropy | 92 |
| 3.4.2. Flory's athermic solution model | 97 |
| 3.4.3. Staverman's corrective factor | 98 |
| 3.5. Combination of the different concepts: the UNIQUAC model | 101 |
| 3.6. The concept of contribution of groups: the UNIFAC model | 107 |
| 3.6.1. The concept of the contribution of groups | 108 |
| 3.6.2. The UNIFAC model | 108 |
| 3.6.3. The modified UNIFAC model (Dortmund) | 114 |
| 3.6.4. Use of the UNIFAC system in the UNIQUAC model | 114 |
| CHAPTER 4. IONIC SOLUTIONS | 117 |
| 4.1. Reference state, unit of composition and activity coefficients of ionic solutions | 119 |
| 4.2. Debye and Hückel's electrostatic model | 121 |
| 4.2.1. Presentation of the problem | 122 |
| 4.2.2. Notations | 123 |
| 4.2.3. Poisson's equation | 124 |
| 4.2.4. Electrical potential due to the ionic atmosphere | 125 |

| | |
|---|------------|
| 4.2.5. Debye and Hückel's hypotheses | 127 |
| 4.2.6. Debye and Hückel's solution for the potential due to the ionic atmosphere | 132 |
| 4.2.7. Charge and radius of the ionic atmosphere of an ion | 134 |
| 4.2.8. Excess Helmholtz energy and excess Gibbs energy due to charges | 136 |
| 4.2.9. Activity coefficients of the ions and mean activity coefficient of the solution | 138 |
| 4.2.10. Self-consistency of Debye and Hückel's model | 141 |
| 4.2.11. Switching from concentrations to molalities | 144 |
| 4.2.12. Debye and Hückel's law: validity and comparison with experimental data | 146 |
| 4.2.13. Debye and Hückel's limit law | 147 |
| 4.2.14. Extensions of Debye and Hückel's law | 148 |
| 4.3. Pitzer's model | 150 |
| 4.4. UNIQUAC model extended to ionic solutions | 155 |
| CHAPTER 5. DETERMINATION OF THE ACTIVITY OF A COMPONENT OF A SOLUTION | 159 |
| 5.1. Calculation of an activity coefficient when we know other coefficients | 160 |
| 5.1.1. Calculation of the activity of a component when we know that of the other components in the solution | 160 |
| 5.1.2. Determination of the activity of a component at one temperature if we know its activity at another temperature | 162 |
| 5.2. Determination of the activity on the basis of the measured vapor pressure | 164 |
| 5.2.1. Measurement by the direct method | 165 |
| 5.2.2. Method using the vaporization constant in reference II | 166 |
| 5.3. Measurement of the activity of the solvent of the basis of the colligative properties | 168 |
| 5.3.1. Use of measuring of the depression of the boiling point – ebullioscopy | 168 |
| 5.3.2. Use of measuring of the depression of the freezing point – cryoscopy | 170 |
| 5.3.3. Use of the measurement of osmotic pressure | 172 |
| 5.4. Measuring the activity on the basis of solubility measurements | 173 |

| | |
|--|-----|
| 5.4.1. Measuring the solubilities in molecular solutions. | 174 |
| 5.4.2. Measuring the solubilities in ionic solutions | 174 |
| 5.5. Measuring the activity by measuring the distribution of a solute between two immiscible solvents | 176 |
| 5.6. Activity in a conductive solution | 176 |
| 5.6.1. Measuring the activity in a strong electrolyte | 176 |
| 5.6.2. Determination of the mean activity of a weak electrolyte on the basis of the dissociation equilibrium | 180 |
| APPENDICES | 181 |
| APPENDIX 1 | 183 |
| APPENDIX 2 | 193 |
| APPENDIX 3 | 207 |
| BIBLIOGRAPHY | 221 |
| INDEX | 225 |